

Clinical Characterization by Principal Component Analysis of Stress Test ECG

Giovanni Bortolan¹, Ivaylo Christov², Iana Simova³, Nikolay Dimitrov⁴, Irena Jekova²,
Vessela Krasteva²

¹Institute of Biomedical Engineering ISIB-CNR, Padova, Italy

²Institute of Biophysics and Biomedical Engineering, Bulg. Acad. of Sci, Sofia, Bulgaria

³Department of Noninvasive Cardiovascular Imaging and Functional Diagnostics,
National Cardiology Hospital, Sofia, Bulgaria

⁴University Hospital "St Ekaterina", Sofia, Bulgaria

Abstract

The aim of the study is to investigate whether and how QRS-complex and T-wave heterogeneity is influenced by different cardiac risk factors and clinical data.

Digital ECG during stress test was acquired in 106 patients (age 63±10 years, 45 males). Two indices obtained by Principal Component Analysis (PCA): complexity (PCA₁) and non-linear components (PCA₂) were used for the analysis of the heterogeneity of the different clinical groups. Mean, max, and standard deviation values were examined in the study.

Significant difference ($p < 0.01$ ÷ 0.05) between PCA₁ of QRS (PCA₁_QRS) was found between subgroups of patients defined according to the presence or absence of angina pectoris, diabetes mellitus, stroke and smokers. Significant difference for PCA₂_QRS was obtained in the presence of angiographically significant coronary artery disease, diabetes mellitus, positive stress test and triglycerides. For the T wave significant difference was found respectively for PCA₁_T in: myocardial infarction, angiographically significant coronary artery disease and gender and for PCA₂_T in: angiographically significant coronary artery disease, percutaneous coronary intervention and gender.

1. Introduction

The principal component analysis (PCA) is a technique that aims to represent a large number of signals by means of a limited number of fundamental values [1]. When applied to digital ECG tracings, the method determines the "principal components" which represent most of the ECG information. The first three eigenvalues of the PCA provide nearly the total energy of the ECG. Since the mathematical procedure calculates fundamental orthogonal components, PCA analysis of the ECG signal

is a modern approach, which can substitute in a certain degree the vectocardiography based on the orthogonal X, Y, and Z leads [1].

The study of the heterogeneity of the ventricular repolarization has been implicated by long in the analysis of the genesis of ventricular arrhythmias [2]. Theoretical and experimental studies suggest that ventricular repolarization occurs in a nonlinear and inhomogeneous fashion. Measures of repolarization that take into account the T-wave complexity using PCA should be a useful surface ECG marker of heterogeneity of repolarization.

Whilst the diagnostic [3-5] and prognostic [2,6] value of PCA of the T wave has been demonstrated, the effect of physiological factors on the QRS and T wave complexity is unknown. The T wave shape or polarity can be influenced by age, sex, heart rate, body position, autonomic activity, respiration, temperature, electrolyte concentration, food and mental activity [7-11]. It is possible that these factors can also affect T wave complexity. PCA has been used for the investigation of the effect of heart rate and body position on QRS and T wave complexity [12] and also for the analysis of the post-extrasystolic changes of the T-wave and QRS complex [13].

The aim of the study is to investigate whether and how QRS-complex and T-wave heterogeneity is influenced by different cardiac risk factors and clinical data.

2. Methods and material

2.1. ECG database

We studied 106 patients: age 63±10 years, 45 males, 39 with diabetes mellitus (DM), 85 with AP, 34 with positive stress test, 18 with a history of myocardial infarction (MI), 48 with angiographically significant coronary artery disease (AS-CAD). Controllable risk

Table 1. Distribution of the cardiac risk factors and clinical variables for the whole group of patients. SD – standard deviation; BMI – body mass index; DM – diabetes mellitus; AH – arterial hypertension; MI – myocardial infarction; AS-CAD – angiographically significant coronary artery disease; PCI – percutaneous coronary intervention; n – number.

Clinical variable	Distribution n = 106
Age – mean ± SD	62.8 ± 10.3
Male – n (%)	45 (42%)
BMI – mean ± SD	28.0 ± 4.3
AH – n (%)	96 (90%)
DM – n (%)	39 (36%)
Dyslipidemia – n (%)	87 (81%)
Total cholesterol (mmol/l) – mean ± SD	5.09 ± 1.1
Triglycerides (mmol/l) – mean ± SD	2 ± 1.8
Family history of CAD – n (%)	11 (10%)
Smokers (present or ex) – n (%)	41 (39%)
Angina pectoris – n (%)	85 (80%)
History of MI – n (%)	18 (17%)
Positive stress ECG test – n (%)	34 (32%)
AS-CAD – n (%)	48 (45%)
PCI – n (%)	40 (37%)
Coronary artery bypass grafting – n (%)	10 (9%)
Stroke	6 (6%)

factors (smoke, high blood pressure, high blood cholesterol, and obesity), not controllable risk factors (gender, age and heredity) and other clinical data for this group of patients are presented in Table 1. Ex smokers are considered individuals who quit from at least 6 months.

Ethics: Signing an informed consent was a prerequisite for inclusion in the study. The study protocol was approved by the local ethical committee and complied with the Declaration of Helsinki.

Patients were included regardless of their sex or age. Exclusion criteria were left ventricular systolic dysfunction with ejection fraction < 40%, haemodynamically significant valvular heart disease, history of ventricular tachycardia, patient unable to perform the stress ECG test or unwilling to sign the informed consent.

Digital 12-lead electrocardiograms (ECG) were acquired during stress ECG test using veloergometer (GE Marquette Stress PC ECG Application) – 2-min stages 25W incremental workload.

The test was considered positive in the setting of ≥ 1 mm horizontal or downward-sloping ST depression 80 msec after J-point.

2.2. Principal component analysis

QRS detection algorithm [14] and automatic

delineation of the QRS and T wave onsets and offsets [15] were performed to bound the segments for the PCA estimation.

Two indices by PCA analysis have been used for this analysis: 1) complexity index, the ratio between the second and the first eigenvalues:

$$PCA_1: \lambda_2/\lambda_1,$$

and 2) non linear components (or relative residuum), the ratio between the sum of all the eigenvalues excluding the first three and the sum of all:

$$PCA_2: (\lambda_4+\lambda_5+\dots+\lambda_n)/\sum_{i=1:n}(\lambda_i).$$

PCA₁ and PCA₂ are applied to the QRS complex and to the T wave. Mean values, max values and standard deviation are the parameters examined in the study.

The parameters based on mean and standard deviation of PCA₁ values are defined by:

$$\text{mean_PCA}_1\text{-QRS} = \text{mean}_i(\text{PCA}_1\text{-QRS}_i)$$

$$\text{mean_PCA}_1\text{-T} = \text{mean}_i(\text{PCA}_1\text{-T}_i)$$

$$\text{std_PCA}_1\text{-QRS} = \text{std}_i(\text{PCA}_1\text{-QRS}_i)$$

$$\text{std_PCA}_1\text{-T} = \text{std}_i(\text{PCA}_1\text{-T}_i)$$

In addition in order to capture short term dynamics, the max value of the mean values of 3 consecutive beats are considered:

$$\text{max_3_PCA}_1\text{-QRS} = \text{max}_i(\text{mean}_{i-1:i+1}(\text{PCA}_1\text{-QRS}_i))$$

$$\text{max_3_PCA}_1\text{-T} = \text{max}_i(\text{mean}_{i-1:i+1}(\text{PCA}_1\text{-T}_i))$$

The PCA₂ parameters are calculated in the same way.

2.3. Statistical analysis

We tested the distribution of different parameters using the Kolmogorov Smirnov test. This analysis showed that the majority of the considered parameters did not present a normal distribution. Consequently, we compared the results of different groups with the non parametric Mann-Whitney U test.

3. Results

Results of PCA₁ and PCA₂ for the QRS complex, reported in Table 2, are showing statistical differences between the CAD/noCAD groups only in std_PCA₂_QRS (p=0.02) and max_3_PCA₂_QRS (p=0.04).

Table 3 reports the results of the same parameters for T wave. It appears that all parameters of the “non linear

Table 2. Results for coronary artery disease CAD/noCAD for the QRS complex.

	CAD	noCAD	p
mean_PCA ₁ _QRS	3.06±2.1	3.14±1.9	-
std_PCA ₁ _QRS	0.63±0.45	0.74±0.41	-
max_3_PCA ₁ _QRS	4.73±2.53	5.00±2.33	-
mean_PCA ₂ _QRS	0.78±0.95	0.86±0.80	-
std_PCA ₂ _QRS	0.78±1.1	1.03±0.83	0.02
max_3_PCA ₂ _QRS	4.29±4.9	6.1±6.6	0.04

Table 3. Results for coronary artery disease CAD/noCAD for T wave.

	CAD	noCAD	p
mean_PCA ₁ _T	1.81±0.96	1.95±0.92	-
std_PCA ₁ _T	1.19±0.49	1.41±0.49	0.013
max_3_PCA ₁ _T	5.12±1.78	5.64±1.62	-
mean_PCA ₂ _T	4.60±3.9	6.75±5.6	0.022
std_PCA ₂ _T	4.61±3.9	6.49±5.2	0.039
max_3_PCA ₂ _T	21.9±17.5	31.2±22.1	0.017

Table 4. Results in the angina pectoris AP/noAP comparison: PCA₁ for QRS and PCA₂ for T wave.

	AP	noAP	p
mean_PCA ₁ _QRS	3.3±1.9	2.1±1.9	0.006
std_PCA ₁ _QRS	0.71±0.40	0.59±0.51	-
max_3_PCA ₁ _QRS	5.1±2.3	3.8±2.5	0.024
mean_PCA ₂ _T	5.4±4.3	7.4±7.3	-
std_PCA ₂ _T	5.2±4.0	7.3±7.0	-
max_3_PCA ₂ _T	25.0±17.2	35.3±30.0	-

components” PCA₂ present a statistical difference between the CAD/noCAD groups, while for the “complexity index” PCA₁ only the standard deviation is presenting a statistical difference (p=0.013).

Table 4 shows the results for PCA₁ for QRS complex and PCA₂ for T wave comparing patients with and without AP.

The coronary artery bypass grafting, total cholesterol, dyslipidemia, arterial hypertension, body mass index, age and family history of CAD among the groups of demographic characteristics and risk factor distribution (presented in Table 1) did not show statistical difference for QRS and T wave.

Table 5 reports a summary of all the comparisons with statistically significant difference. Significant difference between the PCA of the QRS was found between subgroups of patients defined according to the presence or absence of AP (p<0.01 for PCA₁), AS-CAD (p<0.05 for PCA₂), DM (p<0.05 for both PCA₁ and PCA₂), positive stress test (p<0.05 for PCA₂), stroke (p<0.01 for PCA₁), triglycerides > 1.7 (p<0.01 for PCA₂), and present or ex smokers (p<0.01 for PCA₁). The only clinical parameters that are not providing statistically significant difference in the QRS complex analysis are: MI, PCI and gender.

Significant difference between the PCA of the T-waves was present between subject with or without a history of MI (p<0.05 for PCA₁), presence or absence of AS-CAD (p<0.05 for PCA₁ and PCA₂), gender (p<0.01 for PCA₁ and PCA₂), and PCI (p<0.05 for PCA₂). The clinical parameters that are not providing statistically significant difference in the T wave analysis are: AP,

Table 5. Summary of the results for PCA₁ and PCA₂ for QRS and T wave. The most significant differences among mean, std and max_3 are illustrated.

Presence vs absence	QRS complex		T wave	
	PCA ₁	PCA ₂	PCA ₁	PCA ₂
MI	-	-	p<0.05	-
AP	p<0.01	-	-	-
AS-CAD	-	p<0.05	p<0.05	p<0.05
DM	p<0.05	p<0.05	-	-
Pos. Stress Test	-	p<0.05	-	-
Stroke	p<0.01	-	-	-
Gender M	-	-	p<0.01	p<0.01
Triglycerides >1.7	-	p<0.01	-	-
Smokers	p<0.01	-	-	-
PCI	-	-	-	p<0.05

CAD, DM, positive stress test, stroke, triglycerides and smokers.

Decreased PCA values were found in the groups of: MI, AS-CAD, DM, positive stress test, triglycerides, smokers and PCI. Increased PCA values were found in: AP and stroke. In addition, females show higher PCA values than males.

4. Discussion

Different cardiac risk factors and clinical parameters have a distinct influence on the two indices based on the Principal Component Analysis applied on the QRS complexes and T-waves during stress ECG Test.

From clinical perspective it is important to predict which patients are at risk for ventricular arrhythmias and sudden cardiac death in order to adopt primary prophylactic measures. History of MI is a well known risk factor for future ventricular arrhythmias and in the present analysis PCA of the T wave showed a significant difference between groups of patients with or without MI, supporting previous knowledge [16]. The population with a history of MI, however, is not a homogeneous one and maybe PCA has the possibility to distinguish those subjects, who are at a particularly increased risk. This will be evaluated in a further study with prospective follow-up of the selected cohort.

Few comparable studies concerning the influence of demographic and clinical variables on QRS or T wave PCA are present in literature [17-19]. Higher values for PCA of the T wave in patients with CAD has been reported [17]. There is some initial work done on QRS complex PCA in the setting of previous MI [18]. Some authors are analyzing the T wave PCA in the course of acute transmural ischemia, induced by prolonged balloon occlusion during PCI [20].

5. Conclusions

The general conclusion of our study is that the cardiac risk factors and clinical data influence the homogeneity of the QRS-complex and T-wave. The indices of the QRS complex based on PCA analysis showed a higher number of significant differences and proved to be more sensitive to the different clinical conditions.

Acknowledgements

This study was supported by a joint project between the Italian Research Council (Institute of Biomedical Engineering ISIB-CNR) and the Bulgarian Academy of Sciences (Institute of Biophysics and Biomedical Engineering).

References

- [1] Extramiana F, Haggui A, Maison-Blanche P, Dubois R, Takatsuki S, Beaufile P, Leenhardt A. T-wave morphology parameters based on Principle Component Analysis reproducibility and dependance of T offset position. *Annals of Noninvasive Electrocardiology* 2007;12:354-63.
- [2] Okin PM, Devereux RB, Fabsitz RR, Lee ET, Galloway JM, Haward BV. Principal component analysis of the T wave and prediction of cardiovascular mortality in American Indians. The strong heart study. *Circulation* 2002;105:714-9
- [3] Priori S, Mortara D, Napolitano C, Paganini V, Cantù F, Cantù G, Schwartz PJ. Evaluation of the spatial aspect of the T-Wave complexity in the long-QT syndrome. *Circulation* 1997;96:3006-12.
- [4] Badilini F, Fayn J, Maison-Blanche P, Leenhardt A, Forlini MC, Denjoy I, Coumel P, Rubel P. Quantitative aspects of ventricular repolarization: Relationship between three-dimensional T Wave loop morphology and scalar QT dispersion. *Annals of Noninvasive Electrocardiology* 1997;2:146-57.
- [5] De Ambroggi L, Aimè E, Ceriotti C, Rovida M, Negroni S. Mapping of ventricular repolarization potentials in patients with arrhythmogenic right ventricular dysplasia. Principal component analysis of the ST-T waves. *Circulation* 1997;96:4314-8.
- [6] Zabel M, Malik M, Hnatkova K, Papademetriou V, Pittaras A, Fletcher RD, Franz MR. Analysis of T-wave morphology from the 12-Lead electrocardiogram for prediction of long-term prognosis in male US veterans. *Circulation* 2002;105:1066-70.
- [7] Rosen IL, Gardberg M. The effects of nonpathologic factors on the electrocardiogram. *Am Heart J* 1957; 53, 494-504.
- [8] Karjalainen J. Functional and myocarditis-induced T-wave abnormalities. Effect of orthostasis, betablockade, and epinephrine. *Chest* 1983; 83, 868-74.
- [9] Batchvarov VN. Circadian pattern of T wave morphology. In: Malik M, Camm AJ (editors). *Dynamic Electrocardiography*, Blackwell Futura, New York 2004, 358-70.
- [10] Bortolan G, Bressan M, Christov I. Gender and age influences in T-loop morphology, *Computers in Cardiology* 2003;30:513-6.
- [11] Bortolan G, Bressan M, Christov I. Longitudinal modifications of the T-loop morphology. *Computers in Cardiology* 2002;29:685-688.
- [12] Batchvarov VN, Bortolan G, Christov II. Effect of heart rate and body position on the complexity of the QRS and T wave in healthy subjects. *Computers in Cardiology* 2008;35:225-8
- [13] Batchvarov VN, Camm AJ. Post-extrasystolic changes in the complexity of the QRS complex and T wave. *Ann Noninvasive Electrocardiol* 2008;13:421-5.
- [14] Christov II. Real time electrocardiogram QRS detection using combined adaptive threshold. *Biomed Eng Online* 2004;3(28) <http://www.biomedicalengineeringonline.com/content/3/1/28>
- [15] Christov I, Simova I. Q-onset and T-end delineation: Assessment of the performance of an automated method with the use of a reference database. *Phys Meas* 2007;28:213-21.
- [16] Karsikas M, Nojonen K, Huikuri H, Seppanen T. New Vectorcardiographic non-planarity measure of T-wave loop improves separation between healthy subjects and myocardial infarction patients. *Conf Proc IEEE Eng Med Biol Soc* 2009;1754-7.
- [17] Al-Zaiti SS, Runco KN, Carey MG. Increased T wave complexity can indicate subclinical myocardial ischemia in asymptomatic adults. *J Electrocardiol* 2011;44:684-8.
- [18] Donnelly MP, Nugent CD, Finlay DD, Rooney NF, Black ND. Diagnosing old MI by searching for a linear boundary in the space of principal components. *IEEE Trans Inf Technol Biomed* 2006;10:476-83.
- [19] Baglivo FH, Arini PD. Study of T-wave morphology parameters based on Principal Components Analysis during acute myocardial ischemia, XVIII Congreso Argentino de Bioingeniería SABI 2011, http://www.sabi2011.fi.mdp.edu.ar/proceedings/SABI/Pdf/SABI2011_76.pdf

Address for correspondence.

Giovanni Bortolan
Institute of Biomedical Engineering, ISIB -CNR
Corso Stati Uniti, 4
35127 Padova
Italy
Giovanni.Bortolan@isib.cnr.it